

**Universidades Lusíada**

Moreira, Paulo

**Understanding the experience of  
psychopathology after intimate partner violence:  
the role of personality**

<http://hdl.handle.net/11067/5712>

**Metadados**

<b>Data de Publicação</b>	2019
<b>Resumo</b>	Objective(s). To fully understand the dynamics of Intimate Partner Violence (IPV) it is necessary to understand the role of personality. The current understanding of which personality characteristics are associated with IPV victimization is, however, far from comprehensive. Given this gap in the literature, our objective was to examine the associations between the dimensions of the psychobiological model of personality and psychopathological symptoms in women who had experienced IPV. Metho...
<b>Editor</b>	Academic Editor
<b>Palavras Chave</b>	s Intimate partner violence, Victims, Personality, Temperament, Character, Psychopathological symptoms, Women, Domestic violence
<b>Tipo</b>	article
<b>Revisão de Pares</b>	yes
<b>Coleções</b>	[ILID-CIPD] Artigos

Esta página foi gerada automaticamente em 2024-04-26T23:50:47Z com  
informação proveniente do Repositório



# Understanding the experience of psychopathology after intimate partner violence: the role of personality

Paulo A.S. Moreira<sup>1,2</sup>, Márcia Pinto<sup>2,3</sup>, C. Robert Cloninger<sup>4</sup>, Daniela Rodrigues<sup>2,5</sup> and Carlos Fernandes da Silva<sup>6,7</sup>

<sup>1</sup> Instituto de Psicologia e de Ciências da Educação, Universidade Lusíada—Norte (Porto), Porto, Portugal

<sup>2</sup> Centro de Investigação em Psicologia para o Desenvolvimento, CIPD, Porto, Portugal

<sup>3</sup> Centro de Acolhimento Temporário Âncora, Associação para o Desenvolvimento de Rebordosa, Rebordosa, Portugal

<sup>4</sup> School of Medicine, Washington University in St. Louis, St. Louis, MO, United States of America

<sup>5</sup> Estabelecimento Prisional de Santa Cruz do Bispo—Masculino, Direção Geral de Reinserção e Serviços Prisionais, Matosinhos, Portugal

<sup>6</sup> Departamento de Educação e Psicologia, Universidade de Aveiro, Aveiro, Portugal

<sup>7</sup> Center for Health Technology and Services Research, CINTESIS, Porto, Portugal

## ABSTRACT

**Objective(s).** To fully understand the dynamics of Intimate Partner Violence (IPV) it is necessary to understand the role of personality. The current understanding of which personality characteristics are associated with IPV victimization is, however, far from comprehensive. Given this gap in the literature, our objective was to examine the associations between the dimensions of the psychobiological model of personality and psychopathological symptoms in women who had experienced IPV.

**Methods.** Using a case-control design, a group of women who had experienced IPV and who were living in shelters ( $n = 50$ ) were compared to a group of control women who had not experienced IPV ( $n = 50$ ). All women completed the Temperament and Character Inventory—Revised and the Brief Symptom Inventory.

**Results.** Victims of IPV showed significantly higher levels of Harm Avoidance and Self-Transcendence, and lower levels of Reward Dependence and Self-Directedness, than the non-IPV control group. Victims of IPV also reported elevated levels of psychopathological symptoms. Personality dimensions showed a broadly consistent pattern of associations across different psychopathological symptoms. A regression analysis indicated that Novelty Seeking was negatively associated with psychopathological symptoms in victims of IPV, but not significantly associated in non-victims.

**Conclusions.** The study highlights the important role of Harm Avoidance and Self-Directedness for understanding psychopathological symptoms. Novelty Seeking appears to play an important role in the expression of individuals' experiences of IPV. These results have important implications for research and practice, particularly the development and implementation of interventions.

Submitted 13 December 2018

Accepted 20 February 2019

Published 29 March 2019

Corresponding author

Paulo A.S. Moreira,  
paulomoreira@por.ulusiada.pt

Academic editor

Bao-Liang Zhong

Additional Information and  
Declarations can be found on  
page 18

DOI 10.7717/peerj.6647

© Copyright  
2019 Moreira et al.

Distributed under  
Creative Commons CC-BY 4.0

## OPEN ACCESS

**Subjects** Psychiatry and Psychology, Women's Health

**Keywords** Intimate partner violence, Victims, Personality, Temperament, Character, Psychopathological symptoms, Women, Domestic violence

## INTRODUCTION

Violence occurring between family members has been broadly defined as domestic violence ([Goodey, 2005](#)). Intimate Partner Violence (IPV) refers to physical, psychological and/or sexual violence that occurs between intimate partners (including cohabiting or divorced intimate partners, independent of gender; [Niolon et al., 2017](#)). IPV is a universal phenomenon ([Devries et al., 2013](#); [O'Doherty et al., 2014](#)), and studies have revealed high rates of violence against women in particular. Worryingly, a recent survey revealed that one in five women in the European Union has been a victim of physical and/or sexual violence by an intimate partner ([European Union Agency for Fundamental Rights, 2014](#)).

The experience of IPV is associated with negative psychological effects ([Golding, 1999](#)), but the type and severity of these effects may depend somewhat on a victim's personality. The present study aimed to evaluate the associations between the dimensions of the psychobiological model of personality ([Cloninger, 1987](#); [Cloninger, 1994](#)) and psychopathological symptoms in women who had experienced IPV. The following subsections will present past research that describes the psychological impact of IPV and the potential influence of personality, and then introduce the psychobiological model of personality.

### IPV and mental health

Exposure to IPV has a negative impact on multiple domains of an individual's biopsychosocial functioning. Research articles that directly address the associations between IPV and mental health are less common than those that describe the physical consequences of IPV ([Devries et al., 2013](#); [Van Parys et al., 2014](#)). Nonetheless, several influential studies have considered the impact of IPV on mental health. A meta-analysis conducted by [Golding \(1999\)](#) reported that IPV has a significant influence on victims' mental health and is a strong predictor of different types of mental disorders. The negative impact of IPV on victims' psychological functioning has been confirmed more recently by two independent systematic reviews ([Dillon et al., 2013](#); [Lagdon, Armour & Stringer, 2014](#)). Fewer studies, however, have considered how individual differences in personality dimensions serve to buffer or enhance the negative impact of IPV on mental health.

The experience of IPV is a stressor with a strong negative impact on the various biopsychological dimensions of health ([Bonomi et al., 2006](#); [Coker et al., 2002](#); [Dillon et al., 2013](#); [Nurius & Macy, 2010](#); [Renner & Whitney, 2010](#); [Zlotnick, Johnson & Kohn, 2006](#)) and on the functioning of victims' children ([Preto & Moreira, 2012](#)). The high neurophysiological activation associated with abusive relationships ([Okuda et al., 2011](#)) impacts on victims' psychological functioning ([Jarvis, Gordon & Novaco, 2005](#); [Leiner et al., 2008](#)) and thereby constitutes a risk factor for the development of psychopathology ([Golding, 1999](#); [Hathaway et al., 2000](#); [Pico-Alfonso et al., 2006](#)) with high levels of comorbidity ([Lagdon, Armour & Stringer, 2014](#); [Armour & Sleath, 2014](#)). Concordantly, the prevalence of mental disorders in victims of IPV is high ([Lagdon, Armour & Stringer, 2014](#); [Okuda et al., 2011](#); [Sesar, Šimic & Dodaj, 2015](#)). Such disorders include depression ([Beydoun et al., 2012](#); [Sesar, Šimic & Dodaj, 2015](#); [Stein & Kennedy, 2001](#)), eating disorders ([Brady, 2008](#)), substance abuse and addiction ([Humphreys et al., 2005](#); [Simonelli, Pasquali](#)

& De Palo, 2014), anxiety (Pico-Alfonso, 2005; Simmons et al., 2008), and trauma related disorders (Coker et al., 2005). Victims of IPV also present a high prevalence of schizoid personality organizations (Pérez-Testor et al., 2007) and paranoid, borderline and schizotypal personality disorders (Khan, Welch & Zilmer, 1993; Pico-Alfonso, Echeburúa & Martinez, 2008). It is, therefore, of paramount importance to acquire a detailed understanding of the association between IPV and mental health in order to improve the effectiveness of clinical interventions and prognoses of victims.

## Personality and mental health

Personality is a strong predictor of mental health and a causal antecedent of mental disorders (Cloninger, 2004). A small body of research has documented how personality influences the relationship between IPV and psychosocial functioning. As an example, the experience of IPV has been shown to reduce the protective effect of high education on the chances of child maltreatment (Salazar et al., 2014). However, studies describing how personality characteristics are associated with the experience of IPV are scarce. As evidence of this, the systematic review conducted by Lagdon, Armour & Stringer (2014) on the effects of experiencing IPV on mental health did not include personality as a search term, and only one study included in the review described personality-related dimensions (Sharhabani-Arzy, Amir & Swisa, 2005).

Although all women who experience IPV are at increased risk for psychopathology, not all women develop the same symptoms. Moreover, those who present the same symptoms do not necessarily experience them with same severity (Nurius & Macy, 2010). To understand a victim's developmental trajectory toward resiliency/vulnerability to psychopathology after experiencing IPV it is necessary to understand the personality organizations and dynamics involved in biopsychological functioning (Josefsson et al., 2013). Personality dimensions are linked to differences in the risk of experiencing IPV (Loas, Cormier & Perez-Diaz, 2011; Sharhabani-Arzy, Amir & Swisa, 2005). These dimensions also play an important role in a victim's functioning, including engagement in abusive or aggressive situations (Hellmuth & McNulty, 2008), failing to stop abusive relationships (Palker-Correll & Marcus, 2004; Renner & Slack, 2006; Walker, 2009), and identifying and selecting coping strategies (Connor-Smith & Flachsbart, 2007). Personality dimensions also mediate the relationship between coping strategies and psychopathology (Auerbach et al., 2010), predict positive and negative aspects of health (Cloninger & Zohar, 2010; Cloninger, Zohar & Cloninger, 2010; Dyrenforth et al., 2010; Gonçalves & Cloninger, 2010; Josefsson et al., 2011), and are associated with responses to trauma, including the development and recovery of post-traumatic stress disorder (Kunst, 2010; North, Abbacchi & Cloninger, 2012). Considering this evidence, it is clear that obtaining a description of the personality organizations associated with IPV will prove useful for understanding the processes of change underlying the adaptive responses of victims (including those that are spontaneous, and those prompted by interventions).

## The psychobiological model of personality

The few studies to evaluate the personality of IPV victims have typically made use of the Minnesota Multiphasic Personality Inventory-2, the Millon Clinical Multiaxial

Inventory-I and II (MCMI-II-[Millon, 1987](#); [Hart, Dutton & Newlove, 1993](#)), the Personality Assessment Inventory (PAI; [Morey, 1991](#); [Tsang & Stanford, 2007](#)), and the Five-Factor Model ([Hellmuth & McNulty, 2008](#)). Such studies, however, are limited by the fact they tested personality models that do not facilitate an investigation of the psychobiological dynamics involved. Fortunately, the psychobiological model is an integrative approach to understanding the concept of personality as a non-linear developmental process within the individual ([Cloninger & Svrakic, 1997](#); [Cloninger, Svrakic & Przybeck, 1993](#); [Svrakic et al., 1993](#)). According to this model, personality comprises two major dimensions: temperament and character.

Temperament refers to the dispositional and stable tendencies to respond to basic emotional stimuli (danger, novelty and reward) with automatic habitual reactions, including inhibition, activation and behavioral maintenance ([Cloninger, Svrakic & Przybeck, 1993](#)). Four temperament dimensions remain stable throughout an individual's development. Novelty Seeking (NS) describes the tendency to respond intensely to novel stimuli as signals of pleasure. Such responses include an active approach to rewarding stimuli and active avoidance of negative stimuli. Individuals with high NS are exploratory, impulsive, and have a disliking for rules and regulations. Harm Avoidance (HA) describes the propensity to respond intensely to signals of punishment or a frustrating loss of reward by inhibiting behavior. People with high HA tend to be anxious, fearful and shy. Reward Dependence (RD) describes the tendency to respond intensely to signals of reward, especially social approval and attachment cues, and individuals with high RD are sentimental, sociable and friendly in attachments. Finally, Persistence (Ps) describes the propensity to maintain specific behaviors, despite frustration and fatigue, in anticipation of a delayed reward following prior intermittent reinforcement. High Ps is linked to eagerness, ambitiousness and perfectionism ([Cloninger, 1999](#)).

Character refers to the higher-order self-regulatory cognitive processes underlying individual differences in goals, motives, values and standards. Character consists of three dimensions that reflect the intrapersonal, interpersonal and transpersonal self ([Cloninger, Svrakic & Przybeck, 1993](#)). Self-Directedness (SD) refers to the individual differences in the intrapersonal self, who is aware of being an autonomous individual with motives, standards and goals. SD functions as the self-regulatory processes in goal-oriented behaviors, meaning that individuals with high SD are responsible, purposeful, resourceful and self-actualizing. Cooperativeness (C) refers to individual differences in the interpersonal self, who is aware of being a member of a group in which each member has rights and responsibilities. People with high C are tolerant, empathic, helpful, principled and compassionate ([Cloninger, Svrakic & Przybeck, 1993](#)). Self-Transcendence (ST) refers to individual differences in the transpersonal self, who is aware of being a part of a larger whole, such as nature or the universe. People characterized by high ST are easily absorbed in what they love to do and are altruistic and spiritual ([Cloninger, Svrakic & Przybeck, 1993](#)).

The psychobiological model of personality was initially developed for clinical assessment in general and clinical populations, but has since proved to be a valid framework for describing normal and abnormal human personality ([Cloninger & Zohar, 2010](#)). The dimensions of the psychobiological model of personality are causal antecedents of several

indicators of mental health, including psychopathology (both Axis I and Axis II conditions as proposed by the DSM-IV-TR). Recent studies support this model's suitability for describing the personality variables underlying health and all components of wellbeing, as well as ill-being, in adults ([Cloninger & Zohar, 2010](#); [Josefsson et al., 2011](#)) and adolescents ([Moreira et al., 2012](#); [Moreira et al., 2015](#); [Schütz, Archer & Garcia, 2013](#)). Thus, this model is a suitable framework from which to investigate the personality characteristics of IPV victims and their associations with psychopathological symptoms.

With the above evidence considered, the aims of the present study were; (a) to identify the unique personality characteristics of women victims of IPV via a comparison with a control group who had not experienced IPV, and (b) to assess whether the experience of IPV moderates the associations between personality characteristics and psychopathological symptoms.

## MATERIALS & METHODS

### Participants

The study used an unmatched case-control design, allowing for the comparison of cases (women who had experienced IPV) and controls (women who had not experienced IPV). Prior to recruitment, we used a freely available online module for calculating the minimum required sample size for unmatched case-control studies ([Sullivan & Soe, 2007](#)). For this calculation; significance level (alpha) was set at .05; power at 80%; proportion of controls with exposure at 20% (based on [European Union Agency for Fundamental Rights, 2014](#)); ratio of sample size, Controls/Cases at 1:1, and proportion of cases with exposure at 80%. The results indicated that a minimum of 28 cases (12 cases and 14 controls) was required. Based on these results, and the prevalence of IPV victims seeking support, we determined that 50 cases and 50 matched controls would be an acceptable sample size. A consecutive sampling method was then used to recruit 50 women from specialist projects that provide support to victims of IPV. The recruitment process was conducted by a team of social workers, medical professionals, and psychologists, all of whom worked for these projects. Controls (who were women known to this team from the local community) were selected so that both groups would be approximately similar in terms of age and occupational status, although no specific matching method was used. Women were eligible to be a control if they confirmed that they had never been victims of IPV and had no formal or informal records or suspicions of IPV. All participants were from the north of Portugal where, according to a relatively recent EU-wide survey, 19% of ever-partnered women aged 18–74 years report experiencing IPV at least once in their lifetime ([European Union Agency for Fundamental Rights, 2014](#)). This study was approved by an Internal Review Board (CIPD-2009-01) at the Research Center for Positive Development (CIPD), Portugal, and only participants who provided written informed consent were included.

### Instruments

#### Personality

We evaluated personality dimensions using the Temperament and Character Inventory–Revised (TCI-R; [Cloninger, 1999](#)), a self-report instrument with 240 items, each scored



on a 5-point Likert scale from 1 (*definitely false*) to 5 (*definitely true*). This instrument captures the four temperament (NS, HA, RD, and Ps), and three character dimensions (SD, C, and ST) of the psychobiological model of personality. Each of these scales is subdivided into between 3 and 5 subscales that capture different aspects of each personality dimension. The Portuguese version of the TCI-R, which we used in this study, has shown good psychometric properties in the Portuguese population ([Moreira et al., 2017](#)).

### ***Psychopathology symptoms***

To evaluate psychopathology, we used the Portuguese version of the Brief Symptom Inventory (BSI; [Derogatis & Spencer, 1982](#)). This instrument consists of 53 items that assess 9 dimensions of psychopathology symptoms. Items are scored using a 5-point Likert scale ranging from 0 (*never*) to 4 (*many times*). The 9 symptom dimensions are: Somatization, Obsession-Compulsion, Interpersonal Sensitivity, Depression, Anxiety, Hostility, Phobic anxiety, Paranoid ideation and Psychoticism. The Portuguese version of the BSI ([Canavarro, 1999](#)) has been shown to have acceptable psychometric properties, with alpha values that vary between .62 (Psychoticism) and .80 (Somatization).

### **Procedures**

#### ***Data collection procedures***

The psychologists, social workers and medical professionals working for the support projects collaborated in the collection of data for the IPV group. Prior to data collection, the study aims were explained to participants (individually) and informed consents were obtained. Participants in the IPV group completed both questionnaires (paper format) in a technician's office. Participants in the non-IPV completed the questionnaires in an office in the community. For both groups, the order in which we presented the two questionnaires (TCI-R and BSI) was counterbalanced to control for possible order effects. Participants took approximately 1 h to complete the questionnaires.

#### ***Data analysis***

We used statistical methods appropriate for non-matched case-control designs given that we did not use a specific matching procedure to pair cases with controls (instead groups were roughly equated based on age and occupational status). First, group differences based on demographic variables were assessed. For categorical demographic variables—civil status (married, single, divorced or other), occupation status, education level (less than secondary school, secondary school or above secondary school), and violence type (psychological, psychological + physical, or psychological + physical + sexual)—group differences were assessed using Cramer's V. Differences in age were assessed using an independent samples *t*-test. Because cases and controls were statistically different for one of these variables (education level), we tested the main effect of group for each personality dimension and psychopathology symptoms using a series of ANCOVAs. For these analyses, education level was included as a covariate. To assess the associations between dimensions of personality and psychopathological symptoms in victims of IPV we calculated Pearson's correlations between TCI-R and BSI subscales. This correlational analysis was only applied to the IPV group ( $n = 50$ ). We interpreted the effect size of correlation coefficients in accordance with

**Table 1** Demographic characteristics of participants.

		IPV ( <i>n</i> = 50)	Non-IPV ( <i>n</i> = 50)	Total ( <i>n</i> = 100)	IPV vs. Non-IPV comparison
		<i>M</i> ( <i>SE</i> )	<i>M</i> ( <i>SE</i> )	<i>M</i> ( <i>SE</i> )	
Civil status	Age	41.2 (1.85)	37.4 (1.53)	39.3 (1.21)	$t = 1.56, p = .122$
		%	%	%	
	Married	60	56	58	
	Single	26	38	32	Cramer's $V = .169$ , $p = .581$
	Divorced	6	2	4	
Occupation status	Other	8	4	6	
	Homemaker	22	10	16	
	Student	2	8	5	Cramer's $V = .252$ , $p = .095$
	Unemployed	14	6	10	
Education	Other occupation	62	76	69	
	Less than Secondary School	84	56	70	Cramer's $V = .453$ , $p < .001$
	Secondary School	16	10	13	
Violence type	Above Secondary School	0	34	17	
	Psychological	20	–	10	
	Psychological + Physical	62	–	31	–
	Psychological + Physical + Sexual	18	–	9	

the guidelines offered by [Cohen \(1988\)](#). Finally, we conducted a series of linear multiple regression analyses, using the entire sample ( $n = 100$ ), to assess whether the experience of IPV interacts with personality dimensions in explaining variance in total symptoms. For these analyses, personality dimensions and IPV (coded as a dummy variable) were mean centered.

## RESULTS


### Demographic statistics

[Table 1](#) presents the demographic characteristics of the study participants. Participants in the IPV group had significantly lower education levels than participants in the non-IPV group, with none of the IPV group achieving higher than a secondary-school level of education (Cramer's  $V = 0.45, p < .001$ ). There were no significant differences between groups in terms of age, civil status, or occupation status. The majority of the IPV group suffered a combination of physical and psychological violence (62%).

### Differences in personality dimensions

[Figure 1](#) presents personality profiles for the IPV and non-IPV groups. In order to standardize scores across the different personality dimensions, [Figure 1](#) presents  $z$  scores (representing number of standard deviations from the mean). The IPV group was characterized by elevated Harm Avoidance and Self-Transcendence, and lower Novelty Seeking, Reward Dependence, Persistence, Self-Directedness and Cooperativeness compared to the non-IPV group. Next, we conducted a series of ANCOVAs (based on raw scores) to assess the significance of these observed differences after controlling for





**Figure 1** Personality profiles for women victims of IPV (IPV Group) and a control group (non-IPV group). Error bars correspond to  $\pm 1$  standard error. NS, Novelty Seeking; HA, Harm Avoidance; RD, Reward Dependence; Ps, Persistence; SD, Self-Directedness; C, Cooperativeness; ST, Self-Transcendence. Full-size [DOI: 10.7717/peerj.6647/fig-1](https://doi.org/10.7717/peerj.6647/fig-1)


group differences in education (see Table 2). These analyses indicated that the largest observed effect of group was for Self-Directedness: the IPV group had significantly lower Self-Directedness than the non-IPV group ( $F = 13.44$ ,  $p < .001$ ). Women victims of IPV also presented significantly lower levels of Reward Dependence ( $F = 7.52$ ,  $p = .007$ ), and significantly higher levels of Harm Avoidance ( $F = 5.52$ ,  $p = .021$ ) and Self-Transcendence ( $F = 7.90$ ,  $p = .006$ ).

### Differences in psychopathology symptoms

Figure 2 presents the mean summed scores of the BSI symptom subscales for the IPV and non-IPV groups. A series of ANCOVAs (see Table 3) indicated that participants in the IPV group had significantly higher levels of all symptoms. The largest effect of group across all symptoms, after controlling for differences in education, was observed for depression ( $F = 45.43$ ,  $p < .001$ ).

### Correlations between personality dimensions and psychopathological symptoms in victims of IPV

The pattern of correlations across symptom dimensions was broadly consistent for each personality dimension in the IPV group (see Tables 4 and 5). Medium-sized ( $|.50| > r > |.30|$ ; Cohen, 1988) correlations were found between Novelty Seeking and psychoticism ( $r = -.42$ ); between Self-Directedness and hostility ( $r = -.41$ ), paranoid ideation ( $r = -.45$ ) and psychoticism ( $r = -.50$ ); and between Self-Transcendence and interpersonal Sensitivity ( $r = -.43$ ). Harm Avoidance was the only temperament



**Figure 2** Mean average summed scores across BSI symptoms for women victims of IPV (IPV Group) and a control group (Non-IPV Group). Error bars correspond to  $\pm 1$  standard error. Somat., Somatization; Obs., Obsession-Compulsion; Int. Sens., Interpersonal Sensitivity; Dep., Depression; Anx., Anxiety; Host., Hostility; Ph.Anx., Phobic Anxiety; Par.Idea., Paranoid Ideology; Psycho., Psychoticism.

Full-size [DOI: 10.7717/peerj.6647/fig-2](https://doi.org/10.7717/peerj.6647/fig-2)

**Table 2** Personality characteristics of women victims of IPV compared to a control group.

Personality dimension	IPV group (n = 50)		Non-IPV group (n = 50)		ANCOVA: main effect of group controlling for education level				Scale reliability $\alpha$
	M	SE	M	SE	F	p	Partial $\eta^2$	Power	
Novelty Seeking	89.16	1.73	94.16	1.56	0.86	.356	.01	.15	.77
Harm Avoidance*	109.32	2.02	98.80	2.15	5.52	.021	.05	.64	.88
Reward Dependence**	97.38	1.81	105.34	1.31	7.52	.007	.07	.78	.78
Persistence	114.26	2.25	120.40	1.77	1.69	.119	.02	.25	.86
Self-Directedness***	127.14	2.66	144.62	2.58	13.44	<.001	.12	.95	.91
Cooperativeness	132.82	2.04	140.66	1.85	3.53	.063	.04	.46	.86
Self-Transcendence**	82.70	1.61	77.86	1.42	7.90	.006	.08	.80	.78

**Notes.**

Main effect of education level (covariate) not presented in ANCOVA output.

\* $p < .05$ .

\*\* $p < .01$ .

\*\*\* $p < .001$ .

dimension positively correlated with psychopathology symptoms: paranoid ideation ( $r = .46$ ), psychoticism ( $r = .43$ ) and interpersonal sensitivity ( $r = .41$ ).

### The moderating effect of IPV on the association between personality and psychopathological symptoms

A correlational analysis indicated that total symptoms were significantly correlated with education level ( $r = -.25$ ,  $p = .011$ ) and type of IPV ( $r = .54$ ,  $p < .001$ ), and consequently

**Table 3** Psychopathological symptoms in women victims of IPV compared to a control group.

Symptoms	IPV group ( <i>n</i> = 50)		Non-IPV group ( <i>n</i> = 50)		ANCOVA: main effect of group controlling for education level				Scale reliability
	<i>M</i>	<i>SE</i>	<i>M</i>	<i>SE</i>	<i>F</i>	<i>p</i>	Partial $\eta^2$	Power	$\alpha$
Somatization <sup>***</sup>	17.06	0.78	11.74	0.62	19.53	<.001	.17	.99	.84
Obsession <sup>***</sup>	17.64	0.73	12.16	0.51	33.38	<.001	.26	1.00	.84
Interpersonal sensitivity <sup>**</sup>	9.48	0.54	7.60	0.33	7.81	.006	.08	.79	.80
Depression <sup>***</sup>	18.10	0.84	10.42	0.55	45.43	<.001	.32	1.00	.91
Anxiety <sup>***</sup>	16.46	0.80	11.10	0.51	26.06	<.001	.21	1.00	.86
Hostility <sup>***</sup>	11.32	0.55	8.62	0.34	16.64	<.001	.15	.98	.77
Phobic anxiety <sup>**</sup>	10.08	0.65	7.16	0.35	11.96	.001	.11	.93	.80
Paranoid ideation <sup>*</sup>	13.10	0.60	11.40	0.47	4.92	.029	.05	.59	.72
Psychoticism <sup>***</sup>	11.58	0.65	7.56	0.35	29.40	<.001	.23	1.00	.80

**Notes.**

Main effect of education level (covariate) not presented in ANCOVA output.

\**p* < .05.

\*\**p* < .01.

\*\*\**p* < .001.

these variables were included as covariates in the following regression models. Age was not significantly correlated with total symptoms in our sample ( $r = .03$ ), although we also included this variable as a covariate given that prior research has shown clear age effects on psychopathology (e.g., [Erskine et al., 2007](#)).

[Table 6](#) presents the results of seven independent regression analyses. After controlling for the effect of IPV, education level, age, and violence type, Harm Avoidance ( $b = .75$ ,  $p < .001$ ) and Self-Directedness ( $b = -.67$ ,  $p < .001$ ) were significant predictors of total symptoms in their respective models. Crucially, while Novelty Seeking was not a significant predictor of symptoms, it was found to have a significant interaction with IPV ( $b = -1.10$ ,  $p = .034$ ). This interaction is plotted in [Fig. 3](#). (Note that we used z scores for Novelty Seeking and raw scores for total symptoms to graph this significant interaction). The simple slopes for this interaction indicated that there was a non-significant positive relationship between Novelty Seeking and total symptoms in the non-IPV controls,  $b = .25$ , 95% CI  $[-0.51, 1.00]$ , and a significant negative relationship between Novelty Seeking and total symptoms in victims of IPV,  $b = -.85$ , 95% CI  $[-1.53, -0.17]$ . IPV did not interact with the remaining six personality dimensions in the prediction of psychopathology symptoms.

## DISCUSSION

The principle finding of this study was that women victims of IPV had a significantly different character and temperament profile to a group of control women who had reported never experiencing IPV. The results also show that personality dimensions were differentially associated with psychopathology symptoms in victims of IPV, and that the experience of IPV moderated the association between Novelty Seeking and total symptoms. The following discussion will address these findings in turn.

**Table 4** Correlations between temperament dimensions and psychopathological symptoms in women victims of IPV ( $n = 50$ ).

Temperament		Psychopathological symptoms								
Subscales	Scales	Somatization	Obsession	Interpersonal sensitivity	Depression	Anxiety	Hostility	Phobic anxiety	Paranoid ideation	Psychoticism
NS1	<b>Novelty seeking</b>	-.070	-.344*	-.244	-.368*	-.173	-.307*	-.055	-.254	-.422*
	Exploratory excitability	-.161	-.317*	-.220	-.356*	-.192	-.372*	-.191	-.310*	-.389*
	Impulsiveness	-.029	-.223	-.169	-.252	-.034	-.198	.006	-.089	-.240
	Extravagance	-.024	-.216	-.108	-.322*	-.035	-.250	.033	-.182	-.357*
NS2	Disorderliness	.053	-.202	-.196	-.074	-.197	.007	.047	-.072	-.158
NS3	<b>Harm avoidance</b>	.199	.380*	.410*	.302*	.328*	.269	.176	.463*	.425*
NS4	Anticipatory worry	.201	.375*	.431*	.332*	.329*	.373*	.101	.535*	.478*
HA1	Fear of uncertainty	.096	.272	.094	.151	.130	.086	.202	.081	.187
HA2	Shyness	-.005	.088	.319*	.074	.171	.127	-.017	.352*	.166
HA3	Fatigability	.331*	.453*	.378*	.366*	.371*	.188	.312*	.386*	.449*
HA4	<b>Reward dependence</b>	-.139	-.227	-.144	-.131	-.331*	-.315*	-.150	-.389*	-.240
RD1	Sentimentality	-.027	.065	.085	.050	-.047	-.054	.073	-.043	.035
RD2	Openness to communication	-.107	-.255	-.319*	-.141	-.292*	-.247	-.164	-.374*	-.252
RD3	Attachment	-.228	-.393*	-.440*	-.339*	-.418*	-.400*	-.177	-.474*	-.448*
RD4	Dependence	-.010	.050	-.281*	.136	-.084	-.115	-.099	-.084	.097
	<b>Persistence</b>	-.180	-.269	-.168	-.211	-.232	-.111	-.060	-.358*	-.287*
PS1	Resistance to stress	-.195	-.352*	-.202	-.243	-.210	-.161	-.200	-.243	-.277
PS2	Work	-.164	-.187	-.192	-.120	-.172	-.099	-.031	-.313*	-.277
PS3	Ambition	-.059	-.135	-.168	-.127	-.136	-.102	.076	-.366*	-.232
PS4	Perfectionism	-.245	-.266	-.205	-.262	-.330*	.023	-.066	-.301*	-.187

**Notes.**

\* $p < 0.05$ .

Values in bold reflect a medium association,  $r > |.30|$ .

**Table 5** Correlations between character dimensions and psychopathological symptoms in women victims of IPV ( $n = 50$ ).

Character		Psychopathological symptoms								
Subscales	Scales	Somatization	Obsession	Interpersonal sensitivity	Depression	Anxiety	Hostility	Phobic anxiety	Paranoid ideation	Psychoticism
	<b>Self-directedness</b>	-.196	-.377*	-.037	-.380*	-.220	-.410*	-.158	-.446*	-.500*
SD1	Responsibility	-.198	-.456*	-.361*	-.411*	-.258	-.365*	-.196	-.426*	-.533*
SD2	Purposeful	-.162	-.390*	-.295*	-.417*	-.265	-.333*	-.170	-.368*	-.499*
SD3	Resourcefulness	-.157	-.361*	-.302*	-.354*	-.162	-.208	-.164	-.432*	-.345*
SD4	Self-acceptance	-.061	-.005	-.230	-.091	-.030	-.383*	.116	-.307*	-.185
SD5	Congruency	-.160	-.186	-.420*	-.171	-.089	-.270	-.150	-.220	-.295*
	<b>Cooperativeness</b>	-.191	-.061	-.125	-.099	-.135	-.301*	-.096	-.324*	-.149
CO1	Social acceptance	-.136	-.015	-.257	.036	-.076	-.263	-.149	-.084	.000
CO2	Empathy	-.186	-.148	.242	-.154	-.201	-.176	-.005	-.378*	-.223
CO3	Helpfulness	-.100	-.059	-.266	-.055	-.178	-.188	-.003	-.395*	-.161
CO4	Compassion	-.228	-.145	-.040	-.239	-.189	-.429*	-.151	-.383*	-.316*
CO5	Conscience	-.096	.094	-.201	.011	.066	-.075	-.015	-.110	.075
	<b>Self-transcendence</b>	-.126	-.172	-.430*	-.161	-.266	-.080	-.009	-.366*	-.253
ST1	Self-forgetful	-.091	-.258	-.313*	-.196	-.160	-.050	-.073	-.184	-.254
ST2	Transpersonal identification	-.079	-.136	-.134	-.119	-.150	-.174	.115	-.419*	-.298*
ST3	Spiritual acceptance	-.092	.053	-.264	-.010	-.248	.046	-.041	-.188	.024

**Notes.**

\* $p < 0.05$ .

Values in bold reflect a medium association,  $r > |.30|$ .

**Table 6** Summary of seven independent regression analyses testing predictors of total symptoms ( $n = 100$ ).

Model	Predictors	$b$	$p$	Model summary			
				$R^2$	$F$	$p$	Power
NS	NS	−0.30	.245	.36	8.62	<.001	1.00
	IPV	17.07	.251				
	NS × IPV <sup>+</sup>	−1.10	.034				
	Age	−0.07	.797				
	Education	−1.93	.668				
	Violence type	4.4.3	.185				
HA	HA <sup>+</sup>	0.75	<.001	.41	10.82	<.001	1.00
	IPV	13.86	.330				
	HA × IPV	0.46	.225				
	Age	−0.13	.960				
	Education	0.93	.833				
	Violence type	4.02	.208				
RD	RD	−0.28	.318	.35	8.50	<.001	1.00
	IPV	15.53	.302				
	RD × IPV	−1.03	.064				
	Age	−0.06	.819				
	Education	−3.00	.504				
	Violence type	4.47	.182				
Ps	Ps	−0.30	.152	.35	8.21	<.001	1.00
	IPV	15.57	.300				
	Ps × IPV	−0.55	.197				
	Age	−0.13	.631				
	Education	−2.85	.528				
	Violence type	4.63	.170				
SD	SD <sup>+</sup>	−0.67	<.001	.44	12.14	<.001	1.00
	IPV	5.96	.673				
	SD × IPV	−0.25	.95				
	Age	0.02	.943				
	Education	0.37	.930				
	Violence type	4.93	.116				
C	C	−0.29	.190	.33	7.78	<.001	1.00
	IPV	13.69	.373				
	C × IPV	−0.45	.303				
	Age	−0.11	.694				
	Education	−2.95	.529				
	Violence type	4.96	.148				

(continued on next page)



**Table 6** (continued)


Model	Predictors	<i>b</i>	<i>p</i>	Model summary			
				<i>R</i> <sup>2</sup>	<i>F</i>	<i>p</i>	Power
ST	ST	−0.27	.336	.34	7.90	<.001	1.00
	IPV	21.45	.160				
	ST × IPV	−0.83	.134				
	Age	−0.21	.439				
	Education	−3.45	.449				
	Violence type	3.93	.247				

**Notes.**

Constant not included in model summaries.

NS, Novelty seeking; HA, Harm avoidance; RD, Reward dependence; Ps, Persistence; SD, Self-Directedness; CO, Cooperativeness; ST, Self-transcendence; IPV, Intimate Partner Violence.

\**p* < .05.



**Figure 3** Significant interaction effect of Novelty Seeking (NS) and IPV on total symptoms.

Full-size [DOI: 10.7717/peerj.6647/fig-3](https://doi.org/10.7717/peerj.6647/fig-3)

## IPV and personality characteristics

The IPV group had a character configuration of low Self-Directedness, low Cooperativeness, and high Self-Transcendence (a *scT* configuration). In terms of temperament, these women had high levels of Harm Avoidance, and low scores in Novelty Seeking, Reward Dependence and Persistence: an *nHrp* configuration. These personality configurations (Cloninger, Svrakic & Przybeck, 1993) imply that these women were more likely to be immature, disorganized, and schizotypal (associated with *scT* configuration), and to have weak social skills and avoidant behavior (associated with *nHrp* configuration). These character and temperament features are typical of psychological organizations associated with schizotypal personality disorder (Cloninger, Bayon & Svrakic, 1998; Svrakic et al.,

2002), one of the most dominant personality disorders in this population (Pérez-Testor *et al.*, 2007; Pico-Alfonso, Echeburúa & Martínez, 2008).

If the personality characteristics associated with the IPV group are considered as antecedent causes of individual differences (indeed studies have demonstrated personality dimensions have predictive validity in prospective studies; Grucza & Goldberg, 2007), our results imply that women inherently more tolerant of routine and conformity, and less willing to explore new situations are more vulnerable to IPV. Low levels of Novelty Seeking may account for a higher tolerance to repeated patterns of abusive interactions and may partially account for the difficulties many women have in leaving abusive relationships (Cloninger, Svrakic & Przybeck, 1993). The IPV group also showed high levels of Harm Avoidance—indeed significantly higher Harm Avoidance than the non-IPV group—a trait associated with high levels of negative affectivity (Cloninger, Svrakic & Przybeck, 1993). Negative emotions have been shown to interfere with executive functioning (Ochsner *et al.*, 2004) and this may influence the ability of IPV victims to conceive of, plan, and then actively seek new experiences (i.e., new relationships). The significantly lower levels of Reward Dependence found in the IPV group compared to the non-IPV group also imply that these women are dispositionally more tolerant of the lack of warm and secure emotional reward from abusive relationships. These findings are consistent with past evidence about the indicators of the biopsychosocial functioning of victims. Neuroticism, as defined by the Five-Factor model of personality, corresponds to high Harm Avoidance and low Self-Directedness in the psychobiological model (Cloninger, 2006). This pattern of high Harm Avoidance and low Self-Directedness was characteristic of the IPV group in our study. Neuroticism (both in victims and aggressors) is as a major predictor of starting/precipitating the aggressive situations of IPV (Hellmuth & McNulty, 2008).

Thus far, we have discussed the present results in terms of how personality may increase vulnerability to IPV. As with all correlational data, our results might equally demonstrate the association between IPV and stress-related personality changes. In support of this proposal, longitudinal research has shown that the experience of traumatic events in adulthood can cause changes to personality (Josefsson *et al.*, 2013) including increases in neuroticism (high Harm Avoidance and low Self-Directedness) and decreases in agreeableness (Löckenhoff *et al.*, 2009). Consequently, some care should be taken before drawing the conclusion that individuals with certain personality profiles are at increased risk of suffering IPV.

## Personality and psychopathology

Personality influences the nature, intensity and frequency of exposure to stress factors. This interferes with the coping strategies used by individuals in stressful situations (Connor-Smith & Flachsbart, 2007). The selection and use of coping strategies has been shown to be correlated with several Big Five personality dimensions, and in particular, neuroticism has been linked to less adaptive strategies (Connor-Smith & Flachsbart, 2007). The low Novelty Seeking, low Reward Dependence, low Persistence, low Self-Directedness and high Harm Avoidance found in our IPV group (which corresponds closely to neuroticism) therefore implies that victims may not use adequate or adaptive problem-solving and cognitive restructuring strategies (Connor-Smith & Flachsbart, 2007). The elevation of

Self-Transcendence found in victims of IPV is also likely to be related to the coping skills necessary for dealing with daily suffering (Coward & Reed, 1996).

Past studies have demonstrated that personality has an influence on psychopathology via its effect on coping strategies (Auerbach et al., 2010), a finding which is consistent with the elevation of psychopathology in our IPV group. High Harm Avoidance is a strong predictor of the presence and severity of anxiety and mood disorders (Cloninger & Zohar, 2010; Gonçalves & Cloninger, 2010) because it potentiates the impact of negative life events and hinders the development of adaptive coping strategies (Hofmann & Bitran, 2007). Individuals with high Harm Avoidance tend to be highly sensitivity to negative life events and have difficulty dealing with their impact (Cole, Logan & Shannon, 2005). Such characteristics may contribute to the high prevalence of psychopathology in victims of IPV (Howard et al., 2010; Nurius & Macy, 2010; Okuda et al., 2011) and their difficulty in stopping abusive relationships (Palker-Correll & Marcus, 2004; Renner & Slack, 2006). In addition, the character dimensions Self-Directedness and Cooperativeness (which were low in our IPV sample) are predictors of a range of positive indicators of health, including subjective wellbeing (Cloninger & Zohar, 2010; Josefsson et al., 2013). Such associations may help explain the low levels of satisfaction in life (Dyrenforth et al., 2010), low quality of life (Adeodato et al., 2005), and the high psychological distress and suicidality (Leiner et al., 2008) among victims of IPV. Finally, personality dimensions have been shown to predict the development of severe conditions after trauma (such as PTSD; North, Abbacchi & Cloninger, 2012), to mediate the relationship between depression and PTSD (Bargai, Ben-Shakhar & Shalev, 2007), and to predict recovery from PTSD in victims (Kunst, 2010). High levels of Self-Directedness and Cooperativeness are a protective factor against the development of PTSD (North, Abbacchi & Cloninger, 2012). Moreover, a positive affect type of personality (low Harm Avoidance and high Self-Directedness) predicts recovery from PTSD among victims.

The final major finding of the present study, in addition to highlighting the important role of Harm Avoidance and Self-Directedness for understanding psychopathological symptoms, was the significant interaction between Novelty Seeking and IPV. Specifically, high Novelty Seeking in women victims of IPV was linked to fewer total symptoms, while high Novelty Seeking in the control group was related to elevated total symptoms. Novelty Seeking thus appears to plays an important role in the expression of individuals' experiences of IPV via the manifestation of psychopathological symptoms. It is important to highlight that symptoms from affective domains (depression and anxiety), elsewhere referred to as internalizing symptoms, are a prevalent feature in individuals with elevated Harm Avoidance, such is the case in women victims of IPV. High Novelty Seeking refers to a behavioral phenotype of sensitivity to novelty that expresses as higher openness to new experiences and a tendency to externalize internal states. High sensitivity to novelty, and in particularly higher openness to new experience, implies that women with high Novelty Seeking express an intolerance for monotony and may be more easily provoked to a fight or flight response (Cloninger, 1987). Such women may have a stronger drive to explore alternatives to their abusive relationships. Individuals high in Novelty Seeking are also more likely to interact with their environments by externalizing their subjective experiences (both

positive and negative), and this can manifest as impulsive overt expressions of feelings, thoughts and internal states. These impulsive overt expressions can serve as important signals about an individual's subjective experience, and can function as a boundary setter and/or a sign of empowerment within an abusive relationship. Moreover, overt expressions of internal states to individuals outside of the relationship can be an important coping mechanism.

Past research has shown that effective interventions for IPV victims promote the development of internal resources and strengths, particularly personal agency mechanisms and socio-cognitive processes (e.g., [Burnette & Cannon, 2014](#); [Hansen, Eriksen & Elklit, 2014](#); [Hoyeck et al., 2014](#)). Results from studies on change processes (both spontaneous and intervention-promoted) in IPV victims converge on the finding that the psychobiological processes associated with change are well captured by personality dimensions. Processes such as self-esteem and self-efficacy, reflection, awareness, meaning making, or self-identity are components of higher-order dimensions such the individuals' systems of beliefs, standards and goals. These in turn refer to crucial dimensions of personality, as is revealed by their inclusion in many theories of personality. For example, an individual's systems of beliefs, standard and goals refer to the dimensions of character in Cloninger's psychobiological model of personality ([Cloninger, Svrakic & Przybeck, 1993](#)).

## Limitations

Some limitations of the present study require consideration. Firstly, we used a cross-sectional design and were therefore unable to draw any conclusions about how personality might change over time because of IPV. Further, the way we analyzed the associations between personality dimensions and psychopathology (correlational) means that the reader should be wary of inferring causality. Future investigations should use appropriate regression analyses and longitudinal designs if aiming to understand whether personality characteristics serve as antecedents to psychopathology. It should be noted that for such analyses a larger sample size than the one used in this study would be required to achieve sufficient power.

Another limitation to consider is that victims of IPV were recruited from specialist projects. The implication of this aspect of the design is that our sample is unlikely to be fully representative of all women victims of IPV. Women seeking support from specialist projects are likely to represent only the most serious experiences of IPV, and will not include the large number of women who conceal their victimization. Furthermore, women who do reveal their experiences of IPV and seek or accept help from such projects may have specific personality profiles that are not representative of all victims. Consequently, the personality profile for IPV victims identified in our study may correspond to a specific subsample of women victims. Future studies with alternative sampling techniques will be required to verify our findings.

A final limitation of this study is that we did not collect data for variables that might mediate the relationship between high Harm Avoidance, low Self-Directedness and involvement in violence, and thus did not control for them. Future studies should therefore

control variables such as alcohol abuse (*Egan & Hamilton, 2008*), effective problem-solving skills, levels of experienced stress (*Hellmuth & McNulty, 2008*), satisfaction with life (*Dyrenforth et al., 2010*), and history of maltreatment in childhood or characteristics of abuse (*Bensley, Van Eenwyk & Simmons, 2003*).

## CONCLUSIONS

Our study highlights the relevance of personality dimensions for understanding the experience of being a victim of IPV. Harm Avoidance and Self-Directedness are of particular importance for understanding an individual's risk for experiencing psychopathological symptoms. Novelty Seeking appears to play an important role in the expression of individuals' experiences of IPV. These results have important implications for research and practice (especially for the development and implementation of interventions). Specifically, they suggest that the personality of each IPV victim needs to be considered when planning interventions and practices, from crisis interventions to medium- and long-term interventions. Indeed, this corresponds to a major challenge in current mental health services: shifting from an emphasis on symptoms to a person-centered approach to individuals.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work is financed by national funds by FCT—Foundation for Science and Technology, under the Project UID/PSI/04375/2016. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
FCT—Foundation for Science and Technology: Project UID/PSI/04375/2016.

### Competing Interests

C. Robert Cloninger is an Academic Editor for PeerJ.

### Author Contributions

- Paulo A.S. Moreira conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Márcia Pinto performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables.
- C. Robert Cloninger contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Daniela Rodrigues performed the experiments.
- Carlos Fernandes da Silva conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.

## Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Director Board of the Centro de Investigação em Psicologia para o Desenvolvimento (CIPD) approved this research (CIPD-2009-01).

## Data Availability

The following information was supplied regarding data availability:

Data is available at Open Science Framework: <https://osf.io/3y2tk>.

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6647#supplemental-information>.

## REFERENCES

- Adeodato VG, Carvalho RR, De Siqueira VR, De Matos FG. 2005. Quality of life and depression in women abused by their partners. *Revista de Saúde Pública* 39(1):108–113 DOI 10.1590/S0034-89102005000100014.
- Armour C, Sleath E. 2014. Assessing the co-occurrence of intimate partner violence domains across the life-course: relating typologies to mental health. *European Journal of Psychotraumatology* 5(1):Article 24620 DOI 10.3402/ejpt.v5.24620.
- Auerbach RP, Abela JRZ, Zhu X, Yao S. 2010. Understanding the role of coping in the development of depressive symptoms: symptom specificity, gender differences, and cross-cultural applicability. *British Journal of Clinical Psychology* 49(4):47–561.
- Bargai N, Ben-Shakhar G, Shalev AY. 2007. Posttraumatic stress disorder and depression in battered women: the mediating role of learned helplessness. *Journal of Family Violence* 22(5):267–275 DOI 10.1007/s10896-007-9078-y.
- Bensley L, Van Eenwyk J, Simmons WK. 2003. Childhood family violence history and women's risk for intimate partner violence and poor health. *American Journal of Preventive Medicine* 25(1):38–44 DOI 10.1016/S0749-3797(03)00094-1.
- Beydoun HA, Beydoun MA, Kaufman JS, Lo B, Zonderman AB. 2012. Intimate partner violence against adult women and its association with major depressive disorder, depressive symptoms and postpartum depression: a systematic review and meta-analysis. *Social Sciences and Medicine* 75(6):959–975 DOI 10.1016/j.socscimed.2012.04.025.
- Bonomi AE, Thompson RS, Anderson M, Reid RJ, Carrell D, Dimer JA, Rivara FP. 2006. Intimate partner violence and women's physical, mental, and social functioning. *American Journal of Preventive Medicine* 30(6):458–466 DOI 10.1016/j.amepre.2006.01.015.
- Brady SS. 2008. Lifetime family violence exposure is associated with current symptoms of eating disorders among both young men and women. *Journal of Traumatic Stress* 21(3):347–351 DOI 10.1002/jts.20335.



- Burnette CE, Cannon C. 2014.** “It will always continue unless we can change something”: consequences of intimate partner violence for indigenous women, children, and families. *European Journal of Psychotraumatology* 5(1):Article 24585 DOI 10.3402/ejpt.v5.24585.
- Canavarro MC. 1999.** Inventário de sintomas psicopatológicos—BSI. In: Simões MR, Gonçalves M, Almeida LS, eds. *Testes e provas psicológicas em Portugal*. vol. II. Braga: APPORT/SO.
- Cloninger CR. 1987.** A systematic method for clinical description and classification of personality variants. *Archives of General Psychiatry* 44(6):573–588 DOI 10.1001/archpsyc.1987.01800180093014.
- Cloninger CR. 1994.** Temperament and personality. *Current Opinion in Neurobiology* 4(2):266–273 DOI 10.1016/0959-4388(94)90083-3.
- Cloninger CR. 1999.** *The temperament and character inventory—revised*. St. Louis: Center for Psychobiology of Personality, Washington University.
- Cloninger CR. 2004.** *Feeling good: the science of wellbeing*. Washington: Oxford University Press.
- Cloninger CR. 2006.** Personality as a dynamic psychobiological system. In: Widiger TA, Simonsen E, Sirovatka PJ, Regier DA, eds. *Dimensional models of personality disorders: refining the research agenda for DSM-V*. Washington: American Psychiatric Press, 73–76.
- Cloninger CR, Bayon C, Svrakic DM. 1998.** Measurement of temperament and character in mood disorders: a model of fundamental states as personality types. *Journal of Affective Disorders* 51(1):21–32 DOI 10.1016/s0165-0327(98)00153-0.
- Cloninger CR, Svrakic DM. 1997.** Integrative psychobiological approach to psychiatric assessment and treatment. *Psychiatry* 60(2):120–141 DOI 10.1080/00332747.1997.11024793.
- Cloninger CR, Svrakic DM, Przybeck TR. 1993.** A psychobiological model of temperament and character. *Archives of General Psychiatry* 50(12):975–990 DOI 10.1001/archpsyc.1993.01820240059008.
- Cloninger CR, Zohar AH. 2010.** Personality and the perception of health and happiness. *Journal of Affective Disorders* 128(1–2):24–32 DOI 10.1016/j.jad.2010.06.012.
- Cloninger CR, Zohar AH, Cloninger KM. 2010.** Promotion of well-being in person-centered mental health care. *FOCUS* 8(2):165–179.
- Cohen J. 1988.** *Statistical power analysis for the behavioral sciences*. Hillsdale: Erlbaum.
- Coker AL, Davis KE, Arias I, Desai S, Sanderson M, Brandt HM, Smith PH. 2002.** Physical and mental health effects of intimate partner violence for men and women. *American Journal of Preventive Medicine* 23(4):260–268 DOI 10.1016/S0749-3797(02)00514-7.
- Coker AL, Weston R, Creson DL, Justice B, Blakeney P. 2005.** PTSD Symptoms among men and women survivors of intimate partner violence: the role of risk and protective factors. *Violence and Victims* 20(6):625–643 DOI 10.1891/vivi.20.6.625.

- Cole J, Logan T, Shannon L. 2005.** Intimate sexual victimization among women with protective orders: types and associations of physical and mental health problems. *Violence and Victims* **20**(6):695–715 DOI [10.1891/vivi.20.6.695](https://doi.org/10.1891/vivi.20.6.695).
- Connor-Smith JK, Flachsbart C. 2007.** Relations between personality and coping: a meta-analysis. *Journal of Personality and Social Psychology* **93**(6):1080–1107 DOI [10.1037/0022-3514.93.6.1080](https://doi.org/10.1037/0022-3514.93.6.1080).
- Coward DD, Reed PG. 1996.** Self-Transcendence: a resource for healing at the end of life. *Issues in Mental Health Nursing* **17**(3):275–288 DOI [10.3109/01612849609049920](https://doi.org/10.3109/01612849609049920).
- Derogatis LR, Spencer PM. 1982.** *Administration and procedures: BSI. Manual I*. Baltimore: Clinical Psychometric Research.
- Devries KM, Mak JY, Garcia-Moreno C, Petzold M, Child JC, Falder G, Lim S, Bacchus LJ, Engell RL, Rosenfeld L, Pallitto C, Vos T, Abrahams N, Watts CH. 2013.** Global health. The global prevalence of intimate partner violence against women. *Science* **340**(6140):1527–1528 DOI [10.1126/science.1240937](https://doi.org/10.1126/science.1240937).
- Dillon G, Hussain R, Loxton D, Rahman S. 2013.** Mental and physical health and intimate partner violence against women: a review of the literature. *International Journal of Family Medicine* **2013**:Article 313909 DOI [10.1155/2013/313909](https://doi.org/10.1155/2013/313909).
- Dyrenforth PS, Kashy DA, Donnellan MB, Lucas RE. 2010.** Predicting relationship and life satisfaction from personality in nationally representative samples from three countries: the relative importance of actor, partner, and similarity effects. *Journal of Personality and Social Psychology* **99**(4):690–702 DOI [10.1037/a0020385](https://doi.org/10.1037/a0020385).
- Egan V, Hamilton E. 2008.** Personality, mating effort and alcohol-related violence expectancies. *Addiction Research and Theory* **16**(4):369–381 DOI [10.1080/16066350701850717](https://doi.org/10.1080/16066350701850717).
- Erskine JAK, Kvavilashvili L, Conway MA, Myers L. 2007.** The effects of age on psychopathology, well-being and repressive coping. *Aging & Mental Health* **11**(4):394–404 DOI [10.1080/13607860600963737](https://doi.org/10.1080/13607860600963737).
- European Union Agency for Fundamental Rights. 2014.** *Violence against women: An EUwide survey*. Luxembourg: Publications Office of the European Union.
- Golding JM. 1999.** Intimate partner violence as a risk factor for mental disorders: a meta-analysis. *Journal of Family Violence* **14**(2):99–131 DOI [10.1023/A:1022079418229](https://doi.org/10.1023/A:1022079418229).
- Gonçalves DM, Cloninger CR. 2010.** Validation and normative studies of the Brazilian Portuguese and American versions of the Temperament and Character Inventory—Revised (TCI-R). *Journal of Affective Disorders* **124**(1–2):126–133 DOI [10.1016/j.jad.2009.11.007](https://doi.org/10.1016/j.jad.2009.11.007).
- Goodey J. 2005.** *Victims and victimology: research, policy and practice*. Harlow: Pearson Education.
- Gruza RA, Goldberg LR. 2007.** The comparative validity of 11 modern personality inventories: predictions of behavioral acts, informant reports, and clinical indicators. *Journal of Personality Assessment* **89**(2):167–187 DOI [10.1080/00223890701468568](https://doi.org/10.1080/00223890701468568).
- Hansen NB, Eriksen SB, Elklit A. 2014.** Effects of an intervention program for female victims of intimate partner violence on psychological symptoms and perceived

- social support. *European Journal of Psychotraumatology* 5(1):Article 24797 DOI 10.3402/ejpt.v5.24797.
- Hart SD, Dutton DA, Newlove T. 1993. The prevalence of personality disorder among wife assaulters. *Journal of Personality Disorders* 7(4):329–341 DOI 10.1521/pedi.1993.7.4.329.
- Hathaway JE, Mucci LA, Silverman JG, Brooks DR, Mathews R, Pavlos CA. 2000. Health status and health care use of Massachusetts women reporting partner abuse. *American Journal of Preventive Medicine* 19(4):302–307 DOI 10.1016/s0749-3797(00)00236-1.
- Hellmuth JC, McNulty JK. 2008. Neuroticism, marital violence and the moderating role of stress and behavior skills. *Journal of Personality and Social Psychology* 95(1):166–180 DOI 10.1037/0022-3514.95.1.166.
- Hofmann SG, Bitran S. 2007. Sensory-processing sensitivity in social anxiety disorder: relationship to harm avoidance and diagnostic subtypes. *Journal of Anxiety Disorders* 21(7):944–954 DOI 10.1016/j.janxdis.2006.12.003.
- Howard LM, Trevillion K, Khalifeh H, Woodall A, Agnew-Davies R, Feder G. 2010. Domestic violence and severe psychiatric disorders: prevalence and interventions. *Psychological Medicine* 40(6):881–893 DOI 10.1017/S0033291709991589.
- Hoyeck P, Madden K, Freeman C, Scott T, Bhandari M. 2014. Predictors of change in mental health and distress among women attending a women’s shelter. *European Journal of Psychotraumatology* 5:Article 24809 DOI 10.3402/ejpt.v5.24809.
- Humphreys C, Regan L, River D, Thiara RK. 2005. Domestic violence and substance use: tackling complexity. *British Journal of Social Work* 35(8):1303–1320 DOI 10.1007/s10896-005-7800-1.
- Jarvis KL, Gordon EE, Novaco RW. 2005. Psychological distress of children and mothers in domestic violence emergency shelters. *Journal of Family Violence* 20(6):389–402 DOI 10.1007/s10896-005-7800-1.
- Josefsson K, Cloninger CR, Hintsanen M, Jokela M, Pulkki-Råback L, Keltikangas-Järvinen L. 2011. Associations of personality profiles with various aspects of well-being: a population-based study. *Journal of Affective Disorders* 133(1):265–273.
- Josefsson K, Jokela M, Cloninger CR, Hintsanen M, Salo J, Hintsanen T, Pulkki-Råback L, Keltikangas-Järvinen L. 2013. Maturity and change in personality: developmental trends of temperament and character in adulthood. *Development and Psychopathology* 25(3):713–727 DOI 10.1017/S0954579413000126.
- Khan FI, Welch TL, Zilmer EA. 1993. MMPI-2 profiles of battered women in transition. *Journal of Personality Assessment* 60(1):100–111 DOI 10.1007/s10896-005-7800-1.
- Kunst MJJ. 2010. Affective personality type, post-traumatic stress disorder symptom severity and post-traumatic growth in victims of violence. *Stress & Health* 27(1):42–51 DOI 10.1007/s10896-005-7800-1.
- Lagdon S, Armour C, Stringer M. 2014. Adult experience of mental health outcomes as a result of intimate partner violence victimisation: a systematic review. *European Journal of Psychotraumatology* 5(1):Article 24794 DOI 10.3402/ejpt.v5.24794.

- Leiner AM, Compton MT, Houry D, Kaslow NJ. 2008.** Intimate partner violence, psychological distress, and suicidality: a path model using data from African American women seeking care in an urban emergency department. *Journal of Family Violence* 23(6):473–481 DOI 10.1007/s10896-008-9174-7.
- Loas G, Cormier J, Perez-Diaz F. 2011.** Dependent personality disorder and physical abuse. *Psychiatry Research* 185(1–2):167–170.
- Löckenhoff CE, Terracciano A, Patriciu NS, Eaton WW, Costa Jr PT. 2009.** Self-reported extremely adverse life events and longitudinal changes in five-factor model personality traits in an urban sample. *Journal of Traumatic Stress* 22(1):53–59 DOI 10.1002/jts.20385.
- Millon T. 1987.** *Manual for millon clinical multi-axial inventory II (MCMI-II)*. Minneapolis: National Computer Systems.
- Moreira PAS, Cloninger CR, Dinis L, Sá L, Oliveira JT, Dias A, Oliveira J. 2015.** Personality and well-being in adolescents. *Frontiers in Psychology* 5:Article 1494 DOI 10.3389/fpsyg.2014.01494.
- Moreira PAS, Cloninger CR, Rocha MJ, Ferreira N, Oliveira J, Gonçalves DM. 2017.** The psychometric properties of the Portuguese version of the TCI-R. *Psychological Reports* 120(6):1178–1199 DOI 10.1177/0033294117711914.
- Moreira PAS, Oliveira JT, Cloninger KM, Azevedo C, Sousa A, Castro J, Cloninger CR. 2012.** The psychometrics and validity of the junior temperament and character inventory in Portuguese adolescents. *Comprehensive Psychiatry* 53(8):1227–1236 DOI 10.1016/j.comppsy.2012.04.014.
- Morey LC. 1991.** *Personality assessment inventory*. Odessa: Psychological Assessment Resources.
- Niolen PH, Kearns M, Dills J, Rambo K, Irving S, Armstead T, Gilbert L. 2017.** *Preventing intimate partner violence across the lifespan: a technical package of programs, policies, and practices*. Atlanta: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- North CS, Abbacchi A, Cloninger CR. 2012.** Personality and posttraumatic stress disorder among directly exposed survivors of the Oklahoma city bombing. *Comprehensive Psychiatry* 53(1):1–8 DOI 10.1016/j.comppsy.2011.02.005.
- Nurius PS, Macy RC. 2010.** Person-oriented methods in partner violence research: distinct biopsychosocial profiles among battered women. *Journal of Interpersonal Violence* 25(6):1064–1093 DOI 10.1177/0886260509340541.
- Ochsner KN, Ray RD, Cooper JC, Robertson ER, Chopra S, Gabrieli JD, Gross JJ. 2004.** For better or for worse: neural systems supporting the cognitive down- and up-regulation of negative emotion. *Neuroimage* 23:483–499 DOI 10.1016/j.neuroimage.2004.06.030.
- O’Doherty LJ, Taft A, Hegarty K, Ramsay J, Davidson LL, Feder G. 2014.** Screening women for intimate partner violence in healthcare settings: abridged Cochrane systematic review and meta-analysis. *British Medical Journal* 348:Article g2913 DOI 10.1136/bmj.g2913.

- Okuda M, Olfson M, Hasin D, Grant BF, Lin K-H, Blanco C. 2011.** Mental health of victims of intimate partner violence: results from a national epidemiologic survey. *Psychiatric Services* **62**(8):959–962 DOI [10.1176/appi.ps.62.8.959](https://doi.org/10.1176/appi.ps.62.8.959).
- Palker-Correll A, Marcus DK. 2004.** Partner abuse, learned helplessness, and trauma symptoms. *Journal of Social and Clinical Psychology* **23**(4):445–462 DOI [10.1521/jscp.23.4.445.40311](https://doi.org/10.1521/jscp.23.4.445.40311).
- Pérez-Testor C, Castillo JA, Davins M, Salamero M, San-Martino M. 2007.** Personality profiles in a group of battered women: clinical and care implications. *Journal of Family Violence* **22**(2):73–80 DOI [10.1007/s10896-006-9057-8](https://doi.org/10.1007/s10896-006-9057-8).
- Pico-Alfonso MA. 2005.** Psychological intimate partner violence: the major predictor of posttraumatic stress disorder in abused women. *Neuroscience and Biobehavioral Reviews* **29**(1):181–193 DOI [10.1016/j.neubiorev.2004.08.010](https://doi.org/10.1016/j.neubiorev.2004.08.010).
- Pico-Alfonso MA, Echeburúa E, Martinez M. 2008.** Personality disorder symptoms in women as a result of chronic intimate male partner violence. *Journal Family Violence* **23**(7):577–588 DOI [10.1007/s10896-008-9180-9](https://doi.org/10.1007/s10896-008-9180-9).
- Pico-Alfonso M, Garcia-Linares MI, Celda-Navarro N, Blasco-Ros C, Echeburúa E, Martinez M. 2006.** The impact of physical, psychological, and sexual intimate male partner violence on women’s mental health: depressive symptoms, posttraumatic stress disorder, state anxiety, and suicide. *Journal of Women’s Health* **15**(5):599–611 DOI [10.1089/jwh.2006.15.599](https://doi.org/10.1089/jwh.2006.15.599).
- Preto M, Moreira PAS. 2012.** Auto-regulação da aprendizagem em filhos de vítimas de violência doméstica contra mulheres (Self-regulation of learning in children and adolescents whose mothers were victims of domestic violence). *Psicologia: Reflexão e Crítica* **25**(4):730–737 DOI [10.1590/S0102-79722012000400012](https://doi.org/10.1590/S0102-79722012000400012).
- Renner LM, Slack KS. 2006.** Intimate partner violence and child maltreatment: understanding intra—and intergenerational connections. *Child Abuse & Neglect* **30**(6):599–617 DOI [10.1016/j.chiabu.2005.12.005](https://doi.org/10.1016/j.chiabu.2005.12.005).
- Renner LM, Whitney SD. 2010.** Examining symmetry in intimate partner violence among young adults using socio-demographic characteristics. *Journal of Family Violence* **25**(2):91–106 DOI [10.1007/s10896-009-9273-0](https://doi.org/10.1007/s10896-009-9273-0).
- Salazar M, Dahlblom K, Solórzano L, Herrera A. 2014.** Exposure to intimate partner violence reduces the protective effect that women’s high education has on children’s corporal punishment: a population-based study. *Global Health Action* **7**(1):Article 24774 DOI [10.3402/gha.v7.24774](https://doi.org/10.3402/gha.v7.24774).
- Schütz E, Archer T, Garcia D. 2013.** Character profiles and adolescents’ self-reported affect. *Personality and Individual Differences* **54**(7):841–844 DOI [10.1016/j.paid.2012.12.020](https://doi.org/10.1016/j.paid.2012.12.020).
- Sesar K, Šimic N, Dodaj A. 2015.** Differences in symptoms of depression, anxiety and stress between victims and perpetrators of intimate partner violence. *Journal of Sociology and Social Work* **3**(2):63–72 DOI [10.15640/jssw.v3n2a7](https://doi.org/10.15640/jssw.v3n2a7).

- Sharhabani-Arzy R, Amir M, Swisa A. 2005.** Self-criticism, dependency and post-traumatic stress disorder among a female group of help-seeking victims of domestic violence in Israel. *Personality and Individual Differences* **38**(5):1231–1240 DOI [10.1016/j.paid.2004.08.006](https://doi.org/10.1016/j.paid.2004.08.006).
- Simmons AN, Paulus MP, Thorp SR, Matthews SC, Norman SB, Stein MB. 2008.** Functional activation and neural networks in women with posttraumatic stress disorder related to intimate partner violence. *Biological Psychiatry* **64**(8):681–690 DOI [10.1016/j.biopsych.2008.05.027](https://doi.org/10.1016/j.biopsych.2008.05.027).
- Simonelli A, Pasquali CE, De Palo F. 2014.** Intimate partner violence and drug addicted women: from explicative models to gender-oriented treatments. *European Journal of Psychotraumatology* **5**(1):Article 24496 DOI [10.3402/ejpt.v5.24496](https://doi.org/10.3402/ejpt.v5.24496).
- Stein MB, Kennedy C. 2001.** Major depressive and post-traumatic stress disorder comorbidity in female victims of intimate partner violence. *Journal of Affective Disorders* **66**(2–3):133–138 DOI [10.1016/S0165-0327\(00\)00301-3](https://doi.org/10.1016/S0165-0327(00)00301-3).
- Sullivan KM, Soe MM. 2007.** Documentation for sample size for an unmatched case-control study. Available at <http://www.openepi.com/PDFDocs/SSCCDoc.pdf>.
- Svrakic DM, Draganic S, Hill K, Bayon C, Przybeck TR, Cloninger CR. 2002.** Temperament, character, and personality disorders: etiologic, diagnostic, treatment issues. *Acta Psychiatrica Scandinavica* **106**(3):189–195 DOI [10.1034/j.1600-0447.2002.02196.x](https://doi.org/10.1034/j.1600-0447.2002.02196.x).
- Svrakic DM, Whitehead C, Przybeck TR, Cloninger RC. 1993.** Differential diagnosis of personality disorders by the seven-factor temperament model and character. *Archives of General Psychiatry* **50**(12):991–999 DOI [10.1001/archpsyc.1993.01820240075009](https://doi.org/10.1001/archpsyc.1993.01820240075009).
- Tsang JA, Stanford MS. 2007.** Forgiveness for intimate partner violence: the influence of victims and offender variables. *Personality and Individual Differences* **42**(4):653–664 DOI [10.1016/j.paid.2006.08.017](https://doi.org/10.1016/j.paid.2006.08.017).
- Van Parys A, Verhamme A, Temmerman M, Verstraelen H. 2014.** Intimate partner violence and pregnancy: a systematic review of interventions. *PLOS ONE* **9**(1):e85084 DOI [10.1371/journal.pone.0085084](https://doi.org/10.1371/journal.pone.0085084).
- Walker LEA. 2009.** *The battered woman syndrome*. Third edition. New York: Springer Publishing Company.
- Zlotnick C, Johnson DM, Kohn R. 2006.** Intimate partner violence and long-term psychosocial functioning in a national sample of American women. *Journal of Interpersonal Violence* **21**(2):262–275 DOI [10.1177/0886260505282564](https://doi.org/10.1177/0886260505282564).